



Final report

BeefLinks: "DietID" Feedbase mapping to raise productivity of cattle

Project code: P.PSH.1245
Prepared by: Philip Vercoe and Zoey Durmic
The University of Western Australia

Date published: 26 April 2024

PUBLISHED BY
Meat & Livestock Australia Limited
PO Box 1961
NORTH SYDNEY NSW 2059

Meat & Livestock Australia acknowledges the matching funds provided by the Australian Government to support the research and development detailed in this publication.

This publication is published by Meat & Livestock Australia Limited ABN 39 081 678 364 (MLA). Care is taken to ensure the accuracy of the information contained in this publication. However MLA cannot accept responsibility for the accuracy or completeness of the information or opinions contained in the publication. You should make your own enquiries before making decisions concerning your interests. Reproduction in whole or in part of this publication is prohibited without prior written consent of MLA.

Abstract

The composition and nutritional characteristics of WA diverse and variable northern rangeland pastures are not well known, which in turn is a major driver of beef cattle productivity in WA rangelands. This project aimed to identify where and what cattle in WA rangelands graze and how this impacts their productivity. Grazing locations and spatial assessment of the cattle movement were estimated by georeferencing cattle using GPS-enabled devices. The grazing 'signatures' were obtained from plant DNA detected in animal faeces by DNA metabarcoding, and this information was then integrated with vegetation and plant nutritional profiles, to understand what makes up the grazing palate for beef cattle in WA rangelands. All of these inputs (grazing location, plant data and presence in faeces) were then linked to animal productivity parameters (i.e. live weight and live weight gain) towards developing better grazing management practices that improve production, deliver a more consistent supply of animals, and improve rangeland health.

The project was impacted by Covid-19 pandemic, restrictions on regional travel in WA, cyclone, flooding, pastoral access, and staffing issues. However, significant progress was made. In this project, a comprehensive approach was taken - from establishing a list of key plants species to be nutritionally mapped, cattle location, grazing patterns, diet selection to an overall effect on animal productivity has been have developed, applied and validated. A body of knowledge was generated of where and what beef cattle graze, and how it may affect the productivity data in the WA rangelands. It was found that there is a great spectrum of nutritive plant profiles (nutritive values, NV) in WA rangelands. It was also confirmed that faecal plant DNA (fDNA) metabarcoding can be used to obtain more precise information on what animals are actually grazing. It was detected where animals are grazing, and then from the land vegetation profiles and fDNA profiles, grazing plants and grazing palates identified. It was also revealed which attributes (vegetation profiles, NV, diet selection) may potentially improve animal production. This project provided an approach and methodology that can be now applied across different regions, landscapes, plant mosaics, as well as over longer period of time. There was also engagement with the key grower groups and producers in the region.

Table of contents

Abstract
Executive summary
1. Background
2. Objectives
3. Methodology
3.1 Overview
3.2 Identify and engage supply chain stakeholders
3.3 Database on WA rangelands plants
3.4 Nutritive values of rangeland plants
3.5 Cattle movements and grazing distribution
3.6 fDNA metabarcoding
3.7 Animal productivity data
3.8 Data integration and analysis
4. Results
4.1 Experiment 1 – database and nutritive traits of WA rangeland plants
4.2 Experiment 2 - Validation of fDNA metabarcoding in rangeland cattle
4.3 Experiment 3 – Diet selection in northern WA rangelands
4.4 Experiment 4 – Cattle georeferencing, diet and productivity – Station A
5. Conclusion
5.1 Key findings
5.2 Benefits to industry
6. Future research and recommendations
7. References
8. Appendix

Executive summary

1. Background

Cattle and grazing in rangelands

The Western Australian rangelands comprise of northern grasslands (Kimberley and Pilbara) and the southern rangelands, characterised by shrublands (Gascoyne, Murchison, Goldfields–Nullarbor). These are the main beef cattle production regions in the state, with land used for beef production estimated at 33M ha, and over 1 million animals at stocking rates of 1–3 cattle/km². Cattle in the pastoral industry are raised free-range, on native pastures and under natural conditions. The profitability of the northern WA beef industry is therefore driven by the nutritional attributes of the plants and what cattle choose to eat.

However, there is significant lack of information on the diet quality and the grazing profiles and palates. There are also significant gaps in understanding how to best (sustainably) utilize the feedbase when considering feed supply and type, rangeland mosaics and land types, any feed supplement requirement, and overall animal productivity and performance. There is evidence that cattle from WA rangelands entering the domestic and international supply chains are reflecting inadequate access to feed supply, poor in-paddock utilisation of feed and low feed conversion efficiencies. This poses a significant threat to the value of the supply chain. The costs of such production incur inefficiencies and costs manifested in pastoral businesses, at backgrounding or feedlotting, or during processing. A long-standing problem identified by cattle producers in the WA rangelands and in extensive pastoral systems is the lack of knowledge on nutritional status of grazing cattle and how best to influence it, particularly through grazing land management.

In the rangeland setting, the diet is not fixed over time and or to a specific grazing area, which poses a significant issue. As animals are in essence free ranging, their grazing palate changes daily. Animals in these production systems graze across diverse locations, characterised by different vegetation, topography and variable seasonal conditions and pasture quality. While animals are selective in what they graze, it is not known what they choose to eat, and if the plant selections made by individual animals always provides the optimal nutritional return. The nutritional characteristics of the plants are also prone to change, and little is known of these eco-geographical variations that may influence the NV of plants that cattle consume. Consequently, there are large differences in body condition and live weight (LW) and LW gain (LWG) in rangeland cattle of similar ages.

Understanding the factors that drive the efficient animal growth and utilisation of feed in the northern rangelands, is a key factor in the supply chain. The targets also include increased animal performance (feed to meat, fertility, time to turn off or greater efficiencies in feedlots), the ease of logistical management within the north-south supply chain, and reduced carbon footprint of production with concurrent enhanced market demand for products.

Another target is reducing the environmental impact of the beef industry, which include sustainable grazing, rangeland health and reduction of carbon (enteric methane) emissions. Identifying antimethanogenic plants in the grazing palate, as well as optimising animal production can help reduce enteric methane production and intensity from rangeland cattle. Further, cattle can affect rangeland health because certain plant species that are preferred by livestock may decrease through overgrazing. Knowing what animal graze and how to protect and better manage the vital natural resources will also help reduce environmental impact of livestock in the rangelands.

Methods to investigate rangeland grazing, diet selection and quality

Investigations around cattle grazing in rangelands are scarce and are limited to ad-hoc observations and anecdotal evidence of what is eaten by livestock. Current prediction of rangeland condition is based on the assessments made at monitoring sites during inspections. It is still relying on methodology from 1980's, where the assessment of rangeland cattle diet relies on visual rating of ground cover or remotely sensing and interpretation of the perceived grazing plants, what is palatable and what is not (DPIRD, 2019). The faecal near infra-red spectroscopy (F.NIRS) has a value predicting digestibility, crude protein, estimated non-grass component and predicted ADG of the diet of cattle grazing tropical pastures (Dixon & Coates, 2009). While a wide range of tropical grassland diets and animal house experiments were used to develop predictions of F.NIRS, these are not representative of WA rangeland plants.

Whilst these approaches provided useful information, the advances in research methodologies now allow more precise and accurate measurements of cattle grazing. With the development of new methodologies such as the global positioning system (GPS), animal tracking can be done with cattle fitted with the GPS-enabled devices (i.e. ear tags or collars) that can provide a real-time precise location of individual cattle – i.e. 'geo-referencing' (Gwatirisa et al., 2022). In parallel, advancement in molecular techniques have led to an emerging technology – faecal DNA (fDNA) metabarcoding that allows to reconstruct botanical composition of diets by detection of plant DNA in the faeces from free-roaming animals (Kartzinel et al., 2015; Valentini et al., 2009). Advantages of this technique is that it is non-invasive and allow large sample sizes, and determination of diet botanical composition at the species level is quicker given the automation of reading numerous plant fDNA sequences at a single time (Pompanon et al., 2012). In a controlled pen study, it has been shown that it can be used to predict composition of forage-type diets (Scasta et al., 2019).

The fDNA technology can therefore be applied for a more detailed and precise information on rangeland cattle diets. While GPS tracking enables detection of the specific location of the animal to match that with feed found at that location, the fDNA metabarcoding can be representative of what plant species they are consuming. Finally, monitoring and detecting individual cattle liveweight (LW) and liveweight gain (LWG) are important for estimation of nutritional requirements and health management.

The main research questions addressed in this project were – where and what do rangeland cattle eat; and can the grazing, and hence animal productivity and the effect of grazing on environment be improved in rangelands. The attempt was made to integrate these components – GPS positioning, grazing and plant nutritional profile, fDNA profiles and animal growth parameters in order to answer these questions.

Hypothesis

Diet in rangeland cattle is influenced by grazing location and individual preference. The nutritive values of plants will also vary according to location and land type. Animal productivity (measured by LW/LWG) will be related to the profile of plants that they graze, i.e. high performing cattle will readily access plants with high nutritive values compared to low-performing animals that will consume those of lower nutrition value.

2. Objectives

The overall objective of this project was to obtain quantitative data on NV, diet selection and the grazing patterns of the cattle in WA rangelands and to assess the link between diet selection and cattle growth rates.

Specific objectives were:

1. Quantify the diversity and variability of pasture species in the WA rangeland feedbase that supports beef production (literature search and database)
2. Examine animal grazing locations (via animal GPS positioning)
3. Examine NV of plant species that make up northern WA rangelands (via wet chemistry and NIRs spectral analysis)
4. Evaluate diet preference/grazing plant selection (via fDNA metabarcoding),
5. Investigate animal performance (LW/LWG from physical or WOW weighing)
6. Correlate animal diet selection with animal productivity

3. Methodology

3.1 Overview

The methodology comprised of synthesis of existing databases, followed by collection and nutritive analyses of targeted plants; geo-referencing of cattle grazing using GPS devices (Ceres tags or GPS collars); identification of diet selection using fDNA metabarcoding; and productivity data using physical LW measurements.

There was one desktop study, followed by laboratory study and three distinct animal experiments. The desktop study collated information on plants likely to be consumed by rangeland cattle in north and central WA rangelands and developed a database. This information is integrated in the database titled 'CN30 - DietID DATABASE' (Appendix 1). The database evaluated following identifiers: Life form, Soil type/habitat, Interim Biogeographic Regionalisation for Australia (IBRA) Region, Location (station), Desirability in pasture, Grazing value, Palatability, Nutritive value, Medicinal/Toxic to livestock. Plants to progress were selected, collected and examined the NV.

Animal Experiment 1 was conducted in central WA rangelands and aimed to validate fDNA metabarcoding methodology for rangeland cattle by associating the plants present in the landscape with plants present in the faeces. Experiment 2 then examined diet selection in the northern WA rangelands using fDNA metabarcoding. Finally, Experiment 3 attempted to link geo-referencing and diet selection with animal productivity.

Apart from experimental studies, the project also identified and engaged supply chain stakeholders in Kimberley, Pilbara, Gascoyne and Southern pastoral regions. The engagements occurred either directly visiting producer farms, or attending/presenting meetings and conferences.

Methodologies used for distribution and nutritive profiles of grazing plants, animal movements and grazing distribution, diet profile/selection, and animal productivity are described below.

3.2 Database on WA rangelands plants

The approach for creating the database followed some of the principles set and applied in the FFI/MLA Project Enrich for native fodder shrubs (FFI, 2011), as well as those recently outlined for native plants in rehabilitation scenarios for the Pilbara region (Durmic et al., 2021). This was a stepwise process that involved a combination of discussions with leading producers in each region through the relevant regional grower group, pastoralists, researchers and other experts, and a targeted literature review of scientific peer review publications and databases.

Information was obtained from relevant literature published in scientific journals, industry and government reports, herbarium and survey data, online resources, catalogues, published books and encyclopaedias, extension publications, postgraduate theses, as well as any public plant databases that can be accessed. This was combined and further refined in consultations with leading producers in the different regions, other project members, as well as agriculture and livestock experts who have experience in the targeted regions. All information related to the species was compiled within the database, followed by a critical analysis of the data and reporting on the findings. The information was sought on the WA Rangeland plants relevant to their i) habitat, distribution and frequency in the landscapes; ii) biological and agronomical properties of the plants, iii) documented use in the livestock and grazing value; iv) plant nutritive profiles and rumen fermentability data (including effect on methane); v) any reported bioactivity, toxicity or anti-nutritive factors/effects, and any adverse or beneficial effects on animal health.

The information was filtered for grazing potential after a search in the FloraBase (<https://florabase.dpaw.wa.gov.au/>) where parameters 'south-west' and 'found in desert only' were removed. A database was then prepared in Excel format, to enable searching and sorting functions. Within the five themes listed above, separate 'attributes' were assigned, and then each attribute will be populated with a numerical or a descriptive value, or information transformed to common denominators such as to YES/NO (i.e. for desirable in pasture) or LOW/MODERATE/HIGH (i.e. for fodder value). When the information was found only at the genus level, or for another species in the genus, the attribute was assigned to all species in the same genus, but marked in brackets and italics. Once the information was compiled, the data was analysed with scoring and weightings against criteria to separate which species are more likely to provide livestock production benefits as well as forage diversity.

3.3 Nutritive values of rangeland plants

Distribution of grazing plants were done by identifying plants in the field, at specified times and locations. The information was recorded in the Fulcrum platform (<https://www.fulcrumapp.com/>) before collecting voucher specimen if required. Plant samples were collected in this manner: after the plant was selected, catalogued and identified, and voucher specimens (if plants were not identified botanically) were collected where required and identified by botanists at the WA Herbarium. Following this, a sample for analyses was collected. A handful of leaves and small branches was taken by hand, placed in a bag and chilled before being frozen at -20°C. Three field replicates from each plant species were collected as separate, individual plants, 5 m – 100 m apart. The plant samples were then freeze-dried and ground before being submitted for analyses. Plant species were analysed in lab assays, by conventional wet chemistry and NIRS analysis to quantify *in vitro* dry matter digestibility (DMD) and protein (nitrogen), fibre and mineral concentration, as per published protocols (Van Soest et al., 1991).

3.4 Cattle movements and grazing distribution

The locations and grazing patterns were documented by tracking the cattle using 'smart' ear tags fitted with GPS, Ceres Tags (Figure 1a, <https://www.cerestag.com/>) or Vence CattleRider™ (Figure 1b, <https://vence.io/>). Briefly, each animal is fitted with a tag or a collar that communicates with a portal (laptop, mobile) via satellite and sends in real-time the exact GPS location of individual cattle. Ceres Tag can fit in the palm of a hand and weighs 32 g. The technology works using low earth orbit satellite machine to machine (M2M) and Internet of Things (IoT) technology that enables monitoring of animals. Once tagged with the Ceres Tag, user receives continuous data points of their locations which

can be used to make observations about grazing patterns or movements of the cattle. In addition to this, cattle movement data was obtained in Experiment 3 using GPS cattle collars. This was done in conjunction with Beeflinks MLA P.PSH.1306 Virtual Fencing Technology Experiment 2, and the complete detailed there. Briefly, CattleRider™ collars (Vence, US) were used in conjunction to radio telemetry masts. Each collar was fitted with a unique number GPS sensor. Animals were contained for 27 days by virtual fence in a medium-sized paddock before sampling.

Figure 1a. Design of Ceres Tags, size and animal fitted with tag




Figure 1b. Design of CattleRider™ collars fitted on cattle



Mapipedia (<https://mapipedia.com/>) was used for geo-referencing of Ceres ear tag fitted cattle, while animals fitted with Vence CattleRider collars were tracked on HerdManager platform (<https://aus.vence.io/#/main/management>)

3.5 fDNA metabarcoding

Faecal matter collected off the ground (Experiment 1), or from the rectum of individual animal (Experiment 2 and 3) were subjected to fDNA metabarcoding to determine the plant composition of the diet, as per published methods for fDNA metabarcoding (Scasta et al., 2019). Briefly, up to 50 g of faecal material was collected off ground or per rectum in sterile DNA-se free vials and stored at -20°C until analyses. The samples were analysed either in Argaly laboratory, France (<https://www.argaly.com/home>, Experiment 1), or eDNA Frontiers laboratory, Western Australia (Experiment 2 and 3, <http://www.ednafrontiers.com/about/>). The details of their methodologies are listed in Appendix 2.

3.6 Animal productivity data

Animal productivity parameters were based on measuring LW at the beginning and at the end of an extended grazing period and calculating LWG.

3.6 Data integration and analysis

Tools within Mapipedia platforms allowed creating and defining geo-references (i.e. paddocks, landscapes), and the addition of important features such as fences and water points. The movement of cattle based on the GPS data was analysed to produce desired outcomes such as 'heat maps' that show grazing location/movements of individual or groups of animals over time. The diet selection and grazing location was then correlated with animal productivity data for individual animals in the observed herd.

4. Results

4.1 Desktop and laboratory study - developing database and NV of WA rangeland plants

Introduction

Plant species with desirable nutritive profiles can be identified using chemical analyses of plant material for nutritive characteristics and an *in vitro* fermentation technique to examine fermentative properties of the plant, where microbial end-products such as total microbial gas and volatile fatty acid (VFA) production are used as an indirect indication of plant digestibility in the rumen. The same technique can also be used as an indication of the bioactivity of the plant, including likely anti-methanogenic potential. This approach was successfully applied to identify plants with nutritive and anti-methanogenic potential amongst Australian native shrubs (Durmic et al., 2010), alternative forages for dairy cattle in Victoria (Durmic et al., 2016) and beef production systems in northern Australia (Durmic et al., 2017).

The aim of this experiment was to identify plants likely to be consumed by rangeland and to examine their nutritive and fermentative profiles. Our hypothesis was that plant species in the WA Rangelands will vary in their chemical composition.

Materials and methods

Database was compiled based on approach described earlier. Over 300 documents (journals and other publications, databases and reports) were examined, and informal interviews with at least ten pastoralists conducted. Plants were collected from eight stations across the Pilbara, Gascoyne, Murchison and Goldfield regions of Northern Western Australia. The nutritive values were determined by NIRS predictions of diet quality as described earlier.

Results

1. Database

A comprehensive 'CN30-DietID DATABASE' (Appendix 1) was compiled. During the data exploration phase, it became evident that there is information about plants that are grazed by cattle in the region, and engagement with the key grower groups and producers within those regions has been invaluable. The database is now populated with a total of 246 plant species, with 133 having a reported fodder potential.

2. Nutritive values

A total of 10-15 key species from each general region/station were selected and collected (Table 1). Whilst they were collected in similar part of the year, there was diversity in land profiles, soil types

and other attributes. The analysis was done for a total of 108 plant samples, comprising of 92 species (some repeated across stations).

There was a range of nutritive value in plants collected. There were 22 species that had DMD higher than 50 g/100g, and the highest DMD was recorded in species *Calotropis procera*, *Cucumis melo*, *Swainsona formosa*, *Ptilotus exaltatus*, *Crotalaria cunninghamii*, *Eremophila forrestii*, *E. maculata* subsp. *brevifolia*, *Ptilotus polakii* and *Dodonaea lobulata*, and consequently these species also had the highest ME (i.e. 10-12 MJ/kg DM). Over half of the species tested had crude protein (CP) greater than 80 g/kg DM, and those with the highest CP content (i.e. greater than 200 g/kg DM) included *Swainsona Formosa*, *Maireana* spp. and *Enchylaena tomentosa*. There were some marked differences between same plant species collected at different locations (stations). For example, *Triodia pungens* from one station had better NV profile to that from the other station. Similarly, *Setaria dielsii* from one station had greater CP to that from the other, but the greatest variability across nutritive value variables was seen in 'marker' species *Ptilotus obovatus*.

Key finding: The nutritive value for 92 WA rangeland plant species reported. Nearly half of these species may qualify as having good/advantageous nutritive value profile.

Table 1: The nutritive value of plant species collected and analysed in P.PSH.1245

Plant species	Common name	Station	NDF	ADF	DMD	OM	ASH	CP	ME MJ/kg DM
g/100 g DM									
<i>Acacia acuminata</i>	Jam	A	53	37	34	92	8	14	4
<i>Acacia aneura</i>	Mulga	B	53	21	59	92	8	13	9
<i>Acacia bivenosa</i>	Two-nerved wattle	C	23	21	61	87	13	13	9
<i>Acacia colei</i>	Cole's Wattle	D	36	33	44	95	5	12	6
<i>Acacia craspedocarpa</i>	Hop Mulga	B	59	23	60	89	11	15	9
<i>Acacia linophylla</i>	Bowgada	E	60	31	48	94	6	8	7
<i>Acacia pruinocarpa</i>	Black Gidgee	D	36	26	40	92	8	12	5
<i>Acacia pyrifolia</i> DC. var. <i>pyrifolia</i>	Kanji bush	D	52	42	39	96	4	12	5
<i>Acacia tetragonophylla</i>	Kurara	C	31	29	52	95	5	15	7
<i>Acacia tetragonophylla</i>	Kurara	B	50	39	34	95	5	7	5
<i>Acacia tetragonophylla</i>	Kurara	F	37	34	48	95	5	14	7
<i>Acacia victoriae</i>	Bardie bush	C	31	24	53	92	8	16	7
<i>Acacia xiphophylla</i>	Snakewood	C	36	32	56	95	5	13	8
<i>Acacia xiphophylla</i>	Snakewood	A	41	32	46	96	4	11	8
<i>Acacia Citrinoviridis</i>	River Jam	D	36	28	45	96	4	13	6
<i>Acacia colletioides</i>	Wait-a-while	A	35	28	42	95	5	12	6
<i>Aerva javanica</i>	Kapok Bush	G	58	44	52	94	6	8	7
<i>Allocasuarina huegeliana</i>	Sheoak	A	56	29	42	94	7	8	6
<i>Alysicarpus muelleri</i>	-	G	76	46	40	98	2	3	5
<i>Astrebula elemoides</i>	Hoop Mitchel Grass	D	87	54	29	97	3	3	3
<i>Atriplex</i> spp.	Saltbush	E	64	34	44	92	8	13	7
<i>Bothriochloa ewartiana</i>	Desert Bluegrass	F	70	43	49	96	4	7	7
<i>Calotropis procera</i>	Giant Milkweed	G	36	23	79	80	20	15	12
<i>Canthium latifolium</i>	Wild Lemon	E	47	26	51	95	5	10	9
<i>Cassia</i> spp.	Cassia	H	57	21	56	90	10	12	9
<i>Cenchrus setiga</i>	Birdwood Grass	F	57	40	51	97	3	8	7
<i>Cenchrus setiga</i>	Birdwood Grass	G	80	51	36	99	1	2	5

<i>Cenchrus ciliaris</i>	Buffel Grass	C	68	40	44	95	5	5	6
<i>Chrysopogon fallax</i>	Ribbon Grass	F	62	41	46	97	3	6	6
<i>Chrysopogon fallax</i>	Ribbon Grass	G	82	52	31	100	0	4	4
<i>Cleome/Arivela viscosa</i>	Asian Spiderflower	D	53	40	52	89	11	11	7
<i>Clerodendrum tomentosum var. lanceolotum</i>	Pilbara Lolly Tree	D	38	24	65	87	13	17	9
<i>Corchorus incanus subsp. incanus</i>	Jute	G	54	39	42	94	6	8	5
<i>Crotalaria cunninghamii</i>	Green Birdflower	G	29	28	68	92	8	13	10
<i>Cucumis melo</i>	Wild cucumber	F	39	25	71	88	12	17	10
<i>Cymbopogon ambiguus</i>	Lemon Scented Grass	B	76	41	39	90	10	5	5
<i>Dactyloctenium radulans</i>	Button Grass	F	54	31	57	91	9	11	8
<i>Dicanthium sericeum subsp. humilus</i>	Annual Bluegrass	F	58	38	50	94	6	6	7
<i>Dodonaea lobulata</i>	Dead Hop Bush	E	46	22	66	93	7	12	12
<i>Enchylaena tomentosa</i>	Ruby Saltbush	D	34	20	61	83	17	21	9
<i>Eragrostis xerophila</i>	Roebourne Plains Grass	D	68	41	47	97	3	7	6
<i>Eragrotis eriopoda</i>	Woolly butt	A	67	34	46	91	9	6	6
<i>Eremophila cuneifolia</i>	Compact Eremophila	D	42	26	64	92	8	16	9
<i>Eremophila forrestii</i>	Wilcox Bush	E	43	25	67	91	9	15	11
<i>Eremophila fraseri</i>	Turpentine Bush	B	47	26	58	94	6	14	9
<i>Eremophila fraseri</i>	Turpentine Bush	D	41	29	64	87	13	11	9
<i>Eremophila fraseri</i>	Turpentine Bush	D	41	28	62	93	7	12	9
<i>Eremophila glabra</i>	Tar Bush	E	36	26	59	94	6	12	9
<i>Eremophila longifolia</i>	Berrigan	A	41	32	43	93	7	12	7
<i>Eremophila maculata subsp. Brevifolia</i>	Fuscia Bush	C	14	22	67	94	6	14	10
<i>Eremophila oldfieldii</i>	Pixie Bush	E	45	24	58	94	6	12	9
<i>Eremophilla spp.</i>	Eremophila	B	48	29	56	94	6	11	9
<i>Eriochloa procera</i>	Spring Grass	F	54	36	51	94	6	7	7
<i>Evolvulus alsinoides var. decumbens</i>	Tropical Speedwell	D	56	42	57	91	9	9	8
<i>Exocarpos aphyllus</i>	Naked Lady	B	47	30	47	95	6	8	8
<i>Gomphrena canescens</i>	Bachelors Buttons	G	65	46	44	95	5	4	6
<i>Hakea preissii</i>	Needlebush	A	65	38	39	95	5	7	5
<i>Halea lorea</i>	Cork tree	C	62	52	39	100	0	6	5

<i>Indigofera linifolia</i>	Narrow-leafed Indigo	G	38	29	54	93	7	10	8
<i>Ipomoea muelleri</i>	Poison Morning Glory	D	38	29	62	90	10	12	9
<i>Maireana glomerifolia</i>	Ball Leaf Bluebush	B	69	41	35	92	8	8	5
<i>Maireana planifolia</i>	Bluebush	D	38	23	58	84	16	17	8
<i>Maireana pyramidata</i>	Sago Bush	B	58	34	41	92	8	8	6
<i>Maireana spp.</i>	Blue bush	A	31	-7	53	80	20	22	10
<i>Mirbelia platylobioides</i>	Mirbelia	G	45	35	60	94	6	14	9
<i>Panicum effusum</i>	Native Panic Grass, hairy Panic Grass	F	62	39	49	94	6	10	7
<i>Poaceae spp.</i>	Grass	B	81	53	26	92	8	3	3
<i>Portulaca oleracea</i>	Pigweed, Pigface	D	30	29	64	86	14	14	9
<i>Ptilotus aervoides</i>	Mat Mulla Mulla	D	61	40	46	94	6	11	6
<i>Ptilotus auriculifolius</i>	Ear-leaved Mulla Mulla	D	48	35	55	89	11	15	8
<i>Ptilotus clementii</i>	Tassel Top Mulla Mulla	D	58	42	48	90	10	12	7
<i>Ptilotus divaricatus</i>	Climbing Mulla Mulla	A	66	36	35	91	10	10	6
<i>Ptilotus exaltatus</i>	Pink Mulla Mulla	D	45	32	69	81	19	14	10
<i>Ptilotus helipteroides</i>	Hairy Mulla Mulla	D	63	41	46	89	11	10	6
<i>Ptilotus obovatus</i>	Cotton bush	C	63	39	41	92	8	10	5
<i>Ptilotus obovatus</i>	Cotton bush	A	64	35	40	92	8	7	6
<i>Ptilotus obovatus</i>	Cotton Bush	D	59	45	51	90	10	9	7
<i>Ptilotus obovatus</i>	Cotton bush	D	57	40	43	94	6	7	6
<i>Ptilotus polakii</i>	Gascoyne Mulla Mulla	D	50	30	67	91	9	8	10
<i>Ptilotus rotundifolius</i>	Royal Mulla Mulla	D	55	42	41	93	7	11	5
<i>Ptilotus schwartzii</i>	Horse Mulla Mulla	A	61	29	46	92	8	12	7
<i>Rhagodia eremaea</i>	Thorny saltbush	C	44	28	60	94	6	18	9
<i>Rhagodia eremaea</i>	Tall Saltbush	B	71	41	34	92	8	7	6
<i>Rhagodia spp.</i>	Rhagodia	A	65	34	51	91	9	15	8
<i>Rhynchosia minima</i>	Mardie Clover	G	46	35	49	92	8	11	7
<i>Scaevola spinescens</i>	Currant Bush	D	38	28	64	92	8	15	9
<i>Sclerolaena cornishiana</i>	Cartwheel Burr	D	53	32	53	93	7	10	7
<i>Senna artemisioides subsp. oligophylla</i>	Bloodbush	D	29	24	54	95	5	13	8
<i>Senna notabilis</i>	Cockroach Bush	D	37	32	52	89	11	13	7

<i>Setaria dielsii</i>	Diel's Pigeon Grass	D	68	41	49	94	6	7	7
<i>Setaria dielsii</i>	Diel's Pigeon Grass	F	58	38	49	94	6	10	7
<i>Solanum lasiophyllum</i>	Flannel bush	A	70	35	52	92	8	9	7
<i>Solanum lasiophyllum</i>	Flannel bush	D	56	39	54	88	12	13	8
<i>Sporobolus australasicus</i>	Fairy Grass	G	85	54	35	97	3	2	4
<i>Swainsona formosa</i>	Sturt's Desert Pea	D	32	26	71	89	11	22	11
<i>Tecticornia doleiformis</i>	Samphire	B	58	25	44	92	8	12	9
<i>Themeda sp. Hamersley</i> (leaves)	Kangaroo Grass	D	69	39	48	96	4	6	7
<i>Themeda sp. Hamersley</i> (seedhead)	Kangaroo Grass	D	78	46	41	98	2	3	5
<i>Triodia pungens</i>	Soft spinifex	C	66	39	47	94	6	8	6
<i>Triodia pungens</i>	Soft spinifex	G	77	49	42	97	3	4	6
<i>Triodia wiseana</i> (seedhead)	Hard Spinifex	D	93	63	23	100	0	1	2
<i>Urochloa gilesii ssp. occidentalis</i>	Summer Grass	D	72	46	44	93	7	7	6
<i>Urochloa occidentalis</i>	Signal Grass	F	58	37	52	93	7	11	7
<i>Vachellia farnesiana</i>	False Mimosa	D	45	33	61	90	10	15	9

4.2 Experiment 1 - Validation of fDNA metabarcoding in rangeland cattle

Introduction

Rangeland cattle in Australia graze freely and roam on open land in search for pasture of good quality. To date, investigations around cattle grazing in rangelands are scarce and are limited to ad-hoc observations and anecdotal evidence of what is eaten by livestock. Very little investigation was done to determine which plants and what quantities are consumed, and what are nutritive and fermentative properties of plants consumed. The fDNA metabarcoding has been developed for environmental studies (Taberlet et al., 2018; Taberlet et al., 2012) and the methodology has been applied in dietary studies in wild ruminants (Craine et al., 2015; Kartzinel et al., 2015). In recent years, it has been extended to nutritional studies in cattle to quantify geographical patterns of cattle diets in North America (Craine et al., 2016), or where animals were fed known rations of forage plants (Scasta et al., 2019). However, research in free-roaming rangeland cattle in Australia is lacking and validation of the fDNA metabarcoding technique is necessary.

This experiment aimed to compare 'perceived grazing species' in central WA rangelands to what cattle may actually be grazing based on fDNA metabarcoding results.

Materials and methods

Experiment was conducted in Oct 2020 at three stations in WA Central Rangelands. Representative plants were identified within a 1 km radius of water points and based on abundance in the landscape and information from the station managers. Assumption was made that plants around watering point will be representative of plants that animals commonly graze. Plants were identified and catalogued. A total of 27 (9 per station) pooled faecal samples off ground were collected for fDNA analysis and sent to Argaly laboratory in France.

Results

The perceived key grazing species at three stations are listed in Table 2. Using this approach, 39 species were identified.

Table 2: List of perceived grazing species at three stations

Station	Plant label	Species	Latin name
A	C1P1	Wait-a-while	<i>Acacia colletioides</i>
A	C1P10	Unknown perennial grass	(<i>Poaceae</i> spp.) CHAL
A	C1P12	Woolly butt	<i>Eragrostis eriopoda</i>
A	C1P13	Snakewood	<i>Acacia xiphophylla</i>
A	C1P16	Kurara	<i>Acacia tetragonophylla</i>
A	C1P19	Rhagodia	<i>Rhagodia</i> spp.
A	C1P20	Prince of Wales Feather	<i>Ptilotus polystachyus</i>
A	C1P4	Sago Bush	<i>Maireana pyramidata</i>
A	C1P7	Jam	<i>Acacia acuminata</i>
A	C2P1	Mulga	<i>Acacia aneura</i>
A	C2P10	Fine leaf mulga	<i>Acacia assimilis</i>
A	C2P15	Horse mulla mulla	<i>Ptilotus schwartzii</i>
A	C2P18	Sheoak	<i>Allocasuarina huegeliana</i>
A	C2P2	Golden Bluebush	<i>Maireana georgei</i>
A	C2P8	Berrigan	<i>Eremophila longifolia</i>
A	C3P1	Flannel Bush	<i>Solanum lasiophyllum</i>
A	C3P14	Unknown perennial grass	(<i>Poaceae</i> spp.) CHAL
A	C3P16	Unknown perennial grass	(<i>Poaceae</i> spp.) CHAL
A	C3P4	Cotton Bush	<i>Ptilotus obovatus</i>
A	C3P8	Needlebush	<i>Hakea preissii</i>
E	M1P13	Tar Bush Poverty	(<i>Eremophila glabra</i> or <i>E. alternifolia</i>)
E	M1P15	Unknown shrub	tba
E	M1P7	Wilcox Shrub	<i>Eremophila forrestii</i>
E	M2P12	Pixie bush	<i>Eremophila oldfieldii</i>
E	M2P4	Hop mulga	<i>Acacia craspedocarpa</i>
E	M2P5	Bowgada (also known as Wanya)	<i>Acacia ramulosa</i> var. <i>linophylla</i>
E	M2P8	Unknown Poverty Shrub	(<i>Eremophila glabra</i> or <i>E. alternifolia</i>)
E	M3P1	Bead Hop Bush	<i>Dodonaea lobulata</i>
E	M3P14	Wild Lemon	<i>Psyrax latifolia</i>
E	M3P8	Unknown shrub	tba
B	W1P12	Naked Lady	<i>Exocarpos aphyllus</i>
B	W1P19	Samphire	<i>Tecticornia doliiformis</i>
B	W1P22	Unknown Grass	(<i>Poaceae</i> spp.) WEB2
B	W1P27	Unknown Grass 2	(<i>Poaceae</i> spp.) WEB1
B	W1P4	Ball Leaf Bluebush	<i>Maireana glomerifolia</i>
B	W1P7	Unknown tree	tba
B	W2P5	Turpentine Bush	<i>Eremophila fraseri</i>
B	W3P20	Silver cassia	<i>Senna artemisioides</i> subsp. \times <i>artemisioides</i>
B	W3P3	Thorny Saltbush, Tall Saltbush	<i>Rhagodia eremaea</i>

However, the fDNA analyses revealed a total of 76 plant species (Tables 3-5). At Station A, there were 64 plant species, at Station H 51 and at Station E 46; some of which overlapped between the stations, others (i.e. *Sweetia fruticosa*) that were unique to the station. However, upon more detailed analyses, many plants were found that are not (commonly) found in Australia, and only 38 of these are reported in WA Rangelands. There was significant individual variability between plant species (diversity and abundance of fDNA) between individual cattle faecal samples.

Table 3: Diversity and abundance (number of sequence reads) of plant species fDNA in 9 cattle dung specimens collected at Station A. In green – found in Australia

Scientific name	Cattle id								
	C1F1	C1F2	C1F3	C2F1	C2F2	C2F3	C3F1	C3F2	C3F3
<i>Acacia</i> sp.	187			2736	9564	4337	19634	19225	3703
<i>Acacia acuminata</i>			33121			2344		927	
<i>Acacia aneura</i>				9979		4781	3186	1975	1541
<i>Acacia cuspidifolia</i>	6592	31538	500			4932	7022		7849

<i>Acacia tetragonophylla</i>							1442		
<i>Acacieae</i>			2346			2344		927	
<i>Amaranthaceae</i>	505								
<i>Aristida</i>		185							
<i>Asclepiadoideae</i>		869							
<i>Asteraceae</i>		829							321
<i>Atriplex</i>			131			291	505		
<i>Bassia eriophora</i>									318
<i>Brassicaceae</i>							682		
<i>Caesalpinoideae</i>	1152	8427	9813	3574	11114	5714	20554	24194	7207
<i>Cardiochlamyaeae</i>						2429		3638	
<i>Carex</i>					765				
<i>Caryophyllales</i>	5746								
<i>Chenopodiaceae</i>					207		1452		
<i>Chloridoideae</i>									
<i>Citrus</i>	905	12002		4211			929		1738
<i>Dodonaeaeae</i>								1019	
<i>Eleusininae</i>									
<i>Eremophila</i>	4830		5821	3701		8169	12296	9145	2670
<i>Eucalyptus longifolia</i>			696					1019	
<i>eudicotyledons</i>	31857	4767	24531	27732	28242	29123	5729	13766	22930
<i>Euphorbia</i>		3468	2750	707				1389	1917
<i>Euphorbia tirucalli</i>								2691	
<i>Euphorbiaceae</i>		412	194						
<i>Exocarpos sparteus</i>								2691	
<i>Fabaceae</i>		425	33244						224
<i>fabids</i>		829	629	748	1437	1104		259	321
<i>Faidherbia albida</i>		187							
<i>Heliantheae alliance</i>			629	748	779	1104			
<i>Indigofera tinctoria</i>								1207	716
<i>Lamiales</i>			191				587	119	
<i>lamiids</i>	109			103		140	103		
<i>Lessertia frutescens</i>					658				
<i>Macrosolen</i>	275								
<i>Macrosolen tricolor</i>			2056			337	875		
<i>Magnoliophyta</i>		134	690	159	331	262	105	109	372
<i>Maireana</i>	36257	4901	22498	29847	30169	30498	6012	14132	24435
<i>Malvaceae</i>	22150	13117		6738	3185	26005	13952	22155	4472
<i>Malvales</i>		117		4800			1685		
<i>Mesangiospermae</i>	5439		1031	2211	831	2004	278	257	712
<i>mimosoid clade</i>	7557	34022	500	10110	1550	9955	11128	4348	9963
<i>Nepetoideae</i>								1349	
<i>Nyctaginaceae</i>							976		
<i>Nyctanthes arbor-tristis</i>							170		
<i>PACMAD clade</i>							2674		
<i>Papilionoideae</i>		1451							
<i>Pentapetalae</i>	23605	2999		7582	3185	25852	14708	5506	2734
<i>Pentapetalae</i>	6081	898	3927	3598		5507	10461	3924	3327
<i>Poaceae</i>								234	297
<i>Poales</i>								234	297
<i>Proteaceae</i>									
<i>Ptilotus</i>	6251						976		
<i>Rhagodia eremaea</i>			131			291	505		
<i>rosids</i>						153	682	16275	
<i>Sapindaceae</i>			696						
<i>Solanaceae</i>	2273								
<i>Solanum lasiophyllum</i>	2273								
<i>Sweetia fruticosa</i>			293						
<i>Tecticornia lepidosperma</i>					207		1452		

Table 4: Diversity and abundance (number of sequence reads) of plant species fDNA in 9 cattle dung specimens collected at E. In green – found in Australia

Scientific name	Cattle id								
	M1F1	M1F2	M1F3	M2F1	M2F2	M2F3	M3F1	M3F2	M3F3
<i>Acacia sp.</i>	521		7566	13789	49515	22769	14506	14024	5473
<i>Acacia acuminata</i>						4267			5600
<i>Acacia aneura</i>				553	538	132			2800
<i>Acacia cuspidifolia</i>	1212		7564		2139	1422			
<i>Acacieae</i>						805			265
<i>Asclepiadoideae</i>							423		
<i>Asteraceae</i>	111		135						
<i>Atriplex</i>				521		278	788		
<i>Bassia eriophora</i>				123					
<i>Caesalpinioideae</i>	1058		8261	15421	57650	25505	16067	16334	5985
<i>Citrus</i>							3436	33456	8746
<i>Dodonaeaeae</i>						122	446		
<i>Eremophila</i>	8943		699	310	3396	2207	7980	18152	5396
<i>Eucalyptus longifolia</i>						122	446		
<i>eudicotyledons</i>	815		3873	6429	12434	9011	12526	22727	18888
<i>Euphorbia</i>									
<i>Euphorbia tirucalli</i>					1805	1319	4725	2236	15380
<i>Exocarpos sparteus</i>						922			
<i>Fabaceae</i>				295	449	3697			5335
<i>fabids</i>	12611		41684	5882	6689	3627			171
<i>Frankenia</i>	38528	27738	26206	27518		5329	2499		
<i>Gentianales</i>							3436	32736	8746
<i>Hakea preissii</i>					7426	1217			
<i>Heliantheae alliance</i>	11477		41275	5316	6689	3396			
<i>Hypobathrum sp. SH-2010</i>								126	
<i>Lamiales</i>	245				307	171	208	134	240
<i>lamiids</i>								583	
<i>Loranthaeae</i>					155	367			
<i>Magnoliophyta</i>			107	119	699	431	172		361
<i>Maireana</i>	38221	27738	29465	33820	13732	15356	15664	24208	21811
<i>Malvaceae</i>					1106		1251		1456
<i>Mesangiospermae</i>					109				
<i>Mesangiospermae</i>	2155		513	627	2096	692	759	1481	2674
<i>mimosoid clade</i>	1212		7691	1578	9536	3795	1454	2310	2957
<i>Nyctanthes arbor-tristis</i>							108		
<i>Pentapetalae</i>	129		257	118	324		715		
<i>Pentapetalae</i>	7221		845	310	3441	2053	7664	18029	5156
<i>Picea abies</i>					196				
<i>Proteaceae</i>					5818	1217			
<i>Quillaja saponaria</i>									112
<i>Rhagodia eremaea</i>				521		278	788		
<i>rosids</i>	111		281		782		1251		1456

<i>Santalum</i>					1805	397	4832	2236	15682
<i>Solanaceae</i>			895						
<i>Solanum lasiophyllum</i>			895						
<i>Spermatophyta</i>					109				

Table 5: Diversity and abundance (number of sequence reads) of plant species fDNA in 9 cattle dung specimens collected at H. In green – found in Australia

Scientific name	Cattle id								
	W1F1	W1F2	W1F3	W2F1	W2F2	W2F3	W3F1	W3F2	W3F3
<i>Acacia sp.</i>	19876	24235	30021	42705	35534	36818	24510	16517	64007
<i>Acacia acuminata</i>									
<i>Acacia aneura</i>		458	316	390	1573				
<i>Acacia cuspidifolia</i>				3763	352	4977		666	876
<i>Asteraceae</i>		464		178		118			
<i>Atriplex</i>				2065					316
<i>Caesalpinioideae</i>	22407	26648	33201	47478	39004	41997	26742	17975	74199
<i>Caryophyllales</i>					1085	187			
<i>Chenopodiaceae</i>	1548	910	1446	894				130	520
<i>Chloridoideae</i>		142							
<i>Dodonaeaeae</i>									
<i>Eleusininae</i>		10299							
<i>Eremophila</i>									216
<i>Eucalyptus longifolia</i>									
<i>eudicotyledons</i>	1117	4402	717	2400	9300	7294	6805	4577	7188
<i>Euphorbia</i>									
<i>Euphorbia tirucalli</i>	2766	710	633		775	1220	1789		3628
<i>Euphorbiaceae</i>									
<i>Exocarpos sparteus</i>	2766		633		775	1220	1789		3628
<i>Fabaceae</i>	495		684		142				153
<i>fabids</i>	19015	29061	12167	5137	994	20684	13830	13503	2209
<i>Faidherbia albida</i>									
<i>Frankenia</i>						234		339	
<i>Gentianales</i>									
<i>Hakea preissii</i>								211	
<i>Heliantheae alliance</i>	19015	29024	12167	4959	994	20566	13830	13503	2209
<i>Hypobathrum sp. SH-2010</i>									
<i>Indigofera tinctoria</i>									
<i>Lamiales</i>									
<i>lamiids</i>									
<i>Lessertia frutescens</i>									
<i>Loranthaeae</i>									
<i>Macrosolen</i>									
<i>Macrosolen tricolor</i>									216

<i>Magnoliophyta</i>		150		158	452	345	331	239	465
<i>Maireana</i>	1117	4552	1642	2841	10486	8157	8144	5338	8211
<i>Mesangiospermae</i>		11244							
<i>Mesangiospermae</i>	356	495	308	120	389	768	794	497	405
<i>mimosoid clade</i>	1241	1803	1958	8514	5160	8680	1268	1373	10698
<i>Musa</i>				283					
<i>PACMAD clade</i>		803							
<i>Pentapetalae</i>	439	573	1471	292	438	992	1178	437	370
<i>Proteaceae</i>								211	
<i>Ptilotus</i>					1085	187			
<i>Rhagodia eremaea</i>									316
<i>rosids</i>		427							
<i>Salicornia subg. Salicornia</i>	8309	860	2814	580		403	947	2420	
<i>Santalum</i>		710							
<i>Solanaceae</i>	130								
<i>Solanum lasiophyllum</i>	130								
<i>Tecticornia lepidosperma</i>	9857	1770	4260	1474		403	947	2550	520

When comparing perceived grazing species to those found by fDNA metabarcoding, there was more species detected by later. Also, there was discrepancy between the species detected, with only two species were found as total match by both methodologies, and 8 were 'potential/partial match' (Table 6).

Table 6: Comparison between perceived grazing species and those identified by fDNA metabarcoding at three stations. Note – plants not found in Australia removed

Perceived grazing species	Plant fDNA detected
<i>Acacia aneura</i>	
<i>Acacia sp.</i>	
<i>Acacia xiphophylla</i>	<i>Acacia sp.</i>
<i>Acacia acuminata</i>	<i>Acacia sp.</i>
<i>Acacia colletioides</i>	<i>Acacia sp.</i>
<i>Acacia tetragonophylla</i>	<i>Acacia tetragonophylla</i>
<i>Allocasuarina sp.</i>	
<i>Eragrostis eriopoda</i>	<i>Eragrostis sp.</i>
<i>Eremophila longifolia</i>	
<i>Hakea preissii</i>	
<i>Maireana georgei</i>	
<i>Maireana pyramidata</i>	
<i>Poaceae spp.</i>	<i>Poaceae spp.</i>
<i>Ptilotus obovatus</i>	
<i>Ptilotus sp.</i>	
<i>Ptilotus polystachyus</i>	
<i>Rhagodia spp.</i>	
<i>Solanum lasifolium</i>	<i>Solanum lasiophyllum</i>

Agrostidinae
Acacia axillaris
Acacia clelandii
Acacia crassicarpa
Acacia dawsonii
Acacia floribunda
Acacia pubirhachis
Acacia sp.
Acacia victoriae
Andropogoneae
Aristida
Austrostipa
Aveninae
Cassia sp. AMT-2013
Cenchrinae
Cenchrus ciliaris
Cymbopogon caesius
Cymbopogon sp.
Elymus
Enneapogon asperatus
Enteropogon
Eragrostis desertorum
Eragrostis dielsii
Eragrostis kennedyae
Eragrostis lanicaulis
Eragrostis pergracilis
Eragrostis sp.
Eriachne
Eriachne helmsii
Eriachne pulchella
Hordeinae
Hordeum vulgare
Hyparrhenia anamesa
Leymus karelinii
Leymus sp.
Monachather paradoxus
PACMAD clade
Paspalidium geminatum
Pooideae
Setaria
Sorghastrum secundum
Sporobolus actinocladius
Stipeae

*Thyridolepis**Thyridolepis multiculmis**Tripogonella**Triticeae*

Discussion

It was found that cattle ate a greater diversity of species than the ones initially listed, and that species thought to be grazed were not found in the fDNA samples. Previous work using fDNA metabarcoding has shown notable surprises in diet botanical composition of free-roaming herbivores (Craine et al., 2015), and it is likely that cattle in WA Rangelands also have greater plant palate than originally thought. Degradation of fDNA in the sample off-ground, in the environment may limit the interpretation, particularly in hot regions like WA Rangelands. Despite these drawbacks, fDNA still demonstrated the potential to determine plant species grazed and their relative abundance in the diet. It is also possible that the plant identified as key grazing species do not fully represent the diet of the cattle, and that the most desirable plants to cattle would have been over-grazed around the water point and be in low numbers or non-existent at time of sampling. These plants would exist in higher densities with increasing distance from the water point as the cattle move towards the outer regions of their grazing range, so further studies are needed to expand search and identify such plants.

Key finding: The fDNA can detect greater range of grazing species in WA rangelands. Many additional species that have not been previously reported as grazed by cattle have been identified using this approach. Using off-ground faecal samples and foreign lab may limit the accuracy.

4.3 Experiment 2 – Diet selection in northern WA rangelands

Introduction

In Experiment 2, it was confirmed that fDNA metabarcoding may be a useful tool when identifying diet selection of WA Rangeland cattle. While we identified a range of grazing species in Central WA Rangelands, similar information is lacking for northern WA Rangelands. Further, we identified limitation when we used faecal samples off-ground and a laboratory outside of Australia.

The aim of this experiment was to obtain preliminary data in contrasting grazing environment (Central vs North); obtain individual cattle samples for DNA metabarcoding; validate local lab and regional and database for fDNA metabarcoding analysis; compare 'perceived grazing' to what cattle may actually be grazing based on DNA metabarcoding results; obtain some preliminary relationship between individual weights and the individual diet.

Materials and methods

The Experiment 2 was mirroring the work of Experiment 1 in the northern WA rangelands, but trying to overcome these limitations by collecting individual faecal samples obtained from rectum, and using local WA lab and regional plant database for fDNA metabarcoding. In addition, data of cattle movement were obtained using GPS-enabled cattle collars.

The DietID Experiment 3 was conducted in conjunction with the VFT Experiment 2 in P.PSH.1306. The experiment was conducted in May 2021 at Station D in WA North Rangelands, where cattle were grazing in a paddock for 1 month and GPS location tracked by GPS collars. A total of 10 individual faecal samples directly from cow rectum were collected, their start and finish weights and their GPS location.

The fDNA samples were analysed at the eDNA frontiers at Curtin University, (<https://research.curtin.edu.au/scieng/edna-frontiers/>) and using regional plant sequence database (Plastids of the Pilbara, <https://pilbseq.dbca.wa.gov.au/>), in addition to GenBank.

Results

A total of 52 plant species were identified in the faecal sample of 10 cattle at Station D (Table 7). The diet composition in individual cattle varied considerably; for example in cattle #21, *Pluchea* and *Cullen* species seem to be predominant, while cow #79 seem to have more of *Vachellia* in its diet. There was again a discrepancy between what is perceived as main grazing plants to what animals were actually grazing based on fDNA metabarcoding results, and only two species i.e. *Vachellia farnesiana* and *Portulaca oleracea* were identified by both methodologies.

Table 7: Presence and distribution of plant species fDNA (number of sequence reads) in 10 cattle faecal samples collected at Station D

Species	PilbSeq ID	Cattle id									
		1	33	79	47	65	21	57	69	83	85
<i>Pluchea rubelliflora</i>	33794		-	449			16021			84	
<i>Pluchea dentex</i>	33793										
<i>Pluchea indica</i>	NA										
<i>Pluchea dioscoridis</i>	NA										
<i>Pterocaulon pycnostachyum</i>	NA						2670				
<i>Portulaca oleracea</i>	NA		3959			676			1181	57	
<i>Cullen graveolens</i>	33646			90			64483	6285		69	
<i>Cullen pogonocarpum</i>	33624										
<i>Cullen cinereum</i>	22394										
<i>Cullen australasicum</i>	NA										
<i>Cullen graveolens</i>	33646						1536				
<i>Cullen pogonocarpum</i>	33624										
<i>Cullen cinereum</i>	22394										
<i>Cullen australasicum</i>	NA										
<i>Otholobium sericeum</i>	NA										
<i>Rhynchosia minima</i>	22367										
<i>Rhynchosia malacophylla</i>	NA										
<i>Vachellia farnesiana</i>	33637										
<i>Vachellia tortilis</i>	NA			5898	116		46				
<i>Vachellia farnesiana</i> var. <i>pinetorum</i>	NA										
<i>Vachellia nilotica</i>	NA										
<i>Acacia pinetorum</i>	NA										
<i>Sida trichopoda</i>	33661									167	13852
<i>Sida trichopoda</i>	33661	7									
<i>Sida</i> sp. (<i>Shovelanna Hill</i>)	33562			379	730	69	95		3827		14364
<i>Sida</i> sp. (<i>Barlee Range</i>)	33561										

<i>Sida sp. (Articulation below)</i>	33559										
<i>Sida platycalyx</i>	33558										
<i>Sida sp. (dark green fruits)</i>	33557										
<i>Sida fibulifera</i>	33556										
<i>Sida sp. (spiciform panicles)</i>	33555										
<i>Sida sp. (verrucose glands)</i>	33554										
<i>Sida ectogama</i>	33552										
<i>Sida clementii</i>	33551										
<i>Sida sp. (Shovelanna Hill)</i>	33562										
<i>Sida sp. (Barlee Range)</i>	33561										
<i>Sida sp. (Articulation below)</i>	33559										
<i>Sida platycalyx</i>	33558										
<i>Sida sp. (dark green fruits)</i>	33557										
<i>Sida fibulifera</i>	33556										
<i>Sida sp. (spiciform panicles)</i>	33555										
<i>Sida sp. (verrucose glands)</i>	33554										
<i>Sida ectogama</i>	33552										
<i>Sida clementii</i>	33551										
<i>Urochloa pubigera</i>	22446				36						
<i>Urochloa ramosa / Brachiaria ramosa</i>	NA										
<i>Megathyrsus maximus</i>	NA										
<i>Eriochloa villosa</i>	NA										
<i>Eriochloa procera</i>	NA										
<i>Sorghum bicolor subsp. drummondii</i>	NA										


The presence of main species is summarized in the Table 8. It appears that *Eragrostis dielsii* and *Portulaca oleracea* were grazed by all 10 cattle, *Acacia* by 8 cattle, and other species by a smaller number of animals.

Table 8: Summary of key species grazed by individual cattle

Plant species	Cattle id										Present in diet of no of cattle
	1	2	3	4	5	6	7	8	9	10	
<i>Eragrostis dielsii</i>	■	■	■	■	■	■	■	■	■	■	10
<i>Eremophila</i>	■	■									1
<i>Buffel grass</i>				■	■	■					2
<i>Saltbush</i>			■					■			2
<i>Hordeum vulgare</i>					■	■	■				3
<i>Pterocaulon pycnostachyum</i>	■	■	■	■	■	■	■	■	■	■	5
<i>Portulaca oleracea</i>	■	■	■	■	■	■	■	■	■	■	10
<i>Acacia</i>	■	■	■	■	■	■	■	■	■	■	8

Based on this, it may be possible to visualise an ‘average cattle diet’ (Figure 2).

Figure 2. Illustration of an ‘average cattle diet’ based on fDNA metabarcoding results at Station D.



Key findings: We found that cattle ate a greater diversity of species than the ones originally identified as key species for northern WA Rangelands. Using local lab provided more relevant and accurate results.

4.4 Experiment 3 – Cattle georeferencing, diet selection and productivity at Station A

Introduction

In previous two animal experiments, it was revealed that fDNA metabarcoding can be used to reveal the diet selection in rangeland cattle. While these experiments validated the fDNA metabarcoding in WA rangeland cattle, they did not make an attempt to link these findings to grazing location or animal productivity. The objective of this study was to further obtain quantitative data on diet selection and the grazing patterns of the cattle at Station A in the central WA Rangelands, and to assess any possible link between cattle movement, diet selection and cattle growth rates. Specific aims for Experiment 3 align directly with the overall project aims and include:

1. Profile vegetation and land systems
2. Examine cattle location and movements in the landscape over time
3. Identify diet preference and selection
4. Investigate animal performance over time
5. Examine possible links between animal location and diet with animal productivity

Materials and methods

1. Profiling of vegetation and land systems

Vegetation profiles were estimated based on the Windimurra Monitoring Sites reports for site 1-15 provided by the station owner. The land system map was constructed by Contour Environmental and Agricultural Consulting in 2023 (CEAC, 2023). This approach to mapping has been used in all of the regional rangeland surveys in Western Australia, where a land system is defined as 'an area with a recurring pattern of topography, soils and vegetation'. These recurring patterns were mapped using 1:50,000 scale aerial photography or other remotely sensed images and the land systems are then ground-truthed in the field to verify the mapping. Land systems then form the basis of condition assessments and carrying capacity calculations in the rangelands.

2. Cattle movements

Cattle location and movements were determined using geo-referencing via Ceres tags and Mapipedia. The tags were applied to 60 cattle when yarded during the May 2022 muster, and geolocations (latitude and longitude) of each animal were recorded approximately every four hours. The information was automatically downloaded onto Mapipedia platform for routine monitoring of stock movements. The positions of the animals were considered over a window of time as a movement trajectory in space. A density plot (heat map) of these positions for an individual animal, a group of animals, or the entire herd was estimated.

Diet preference and selection

This was assessed using land system approach and fDNA metabarcoding.

Firstly, the Mapipedia heat map of cattle locations were overlayed onto the land systems map to show land system preferences of each Ceres-tagged animal to determine the grazing behaviour over a 12-month period. Water points were added to determine the likely influence on cattle movements. Preferences for land systems were determined.

Faecal samples were taken from the 26 individual Ceres-tagged animals in April/May 2023 muster and subjected to fDNA metabarcoding at eDNA frontiers lab, as described earlier.

3. Animal performance

Growth performance of cattle was assessed using measuring LW at muster in 2022 and 2023, and calculating LWG. Cattle (n=60) were weighed at mustering in May 2022 and again in late April/early May 2023. Average daily liveweight change (LWG) was used to indicate animal performance over the period of 12 months. At the 2023 muster, 26 of the Ceres-tagged cattle were identified and sampled. The remaining animals were located on Mapipedia, but they could not be yarded during the muster. Animal performance was then classified according to daily weight change to high (> 0.400 kg/d), medium (0.250 – 0.400 kg/d), low (0.100 -0.250 kg/d) and very low (<0.100 kg/d).

4. Correlating animal movement and diet with animal productivity

The 'heat map' of geolocation densities of all Ceres-tagged cattle was refined for cattle with growth performance data (n=26). These geolocations were overlayed onto the land systems map to determine the preferences of land systems for the various growth performance classes.

Results

1. Profiling of vegetation and land systems

There were 37 plant species detected across 15 monitoring sites, with most common and numerous being *Maireana pyramidata*, *Ptilotus obovatus*, *Soliva sessilis* and *Triptria* (Table 9).

Table 9: Vegetation profiles (numbers of plants recorded) at Windimurra Monitoring Sites, Station A Nov 2020

Latin name	Common name	Site number														
		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
<i>Acacia acuminata</i>	Jam tree										1					1
<i>Acacia sp.</i>	Fine Leaf Mulga	1									5	2	5			
<i>Acacia aneura</i>	Mulga			2						2					2	6
<i>Acacia craspedocarpa</i>	Broad Leaf Mulga										1		1		3	
<i>Acacia papyrocarpa</i>	Western Myall		1					1								
<i>Acacia ramulosa</i> var. <i>linophylla</i>	Bowgata			3									1		2	
<i>Acacia stellaticeps</i>	Poverty Bush					1				2			2		1	
<i>Acacia tetragonophylla</i>	Kurara		1		2								2	1	1	2
<i>Agonis flexuosa</i>	Native Willow													1		
<i>Asphodelus fistulosus</i>	Wild Onion												1			
<i>Atriplex bunburyana</i>	Silver Saltbush					5										
<i>Brachychiton populneus</i>	Kurrajong															1
<i>Cratystylis subspinescens</i>	Sage				1											
<i>Enchylaena tomentosa</i>	Ruby Saltbush							1				1				1
<i>Eremophila fraseri</i>	Turpentine Bush			1							3					
<i>Eremophila scoparia</i>	Mulga broombush												1			
<i>Eremophila sp.</i>	Eremophila sp.															1
<i>Frankenia sp.</i>	Frankenia					3										
<i>Maireana pyramidata</i>	Sago Bush	13	23		5	8	4	17	12	9	4	20		18		
<i>Maireana sedifolia</i>	Blue Bush			1												
<i>Ptilotus obovatus</i>	Cotton Bush	1	1	4						10	22			1		25
<i>Ptilotus sp.?</i>	Gascoyne Mulla						7									
<i>Ptilotus sp.?</i>	Wire Mulla			3												
<i>Rhagodia eremaea</i>	Tall Saltbush				1						1	1		2		
<i>Salsola australis</i>	Tumbleweed										1					
<i>Scaevola spinescens</i>	Currant Bushes				2											
<i>Senna artemisioides</i> subsp. <i>x artemisioides</i>	Banana Leaf Cassia					1										
<i>Senna artemisioides</i> subsp. <i>x coriacea</i>	Desert Cassia								9							
<i>Senna sp.</i>	Green Cassia				1											
<i>Senna sp.</i>	White Cassia								2							
<i>Sida calyxhymenia</i>	Tall Sida		4							13	12					
<i>Solanum lasiophyllum</i>	Flannel Bush	4	1			2				1	1					2

<i>Solanum orbiculatum</i>	Wild Tomato							1				
<i>Soliva sessilis</i>	Bindi	5	3				13	100	9		3	
<i>Tecticornia lepidosperma</i>	Samphire						1					
unknown	Triptr		6	2	26	10	5	2	18	4	2	1
unknown	Wild Lemon										1	

There was a total of 28 different land systems identified at Station A.

5. Cattle location and movements

Examining cattle geo-referencing data revealed that cattle had preferred areas in the landscape where the animals spend most of their time.

6. Diet preference/selection

Diet preference

Only 7 land systems were found to be preferably grazed by cattle. The profiles of preferred grazing land systems are given in Table 10.

Table 10: Description of land systems preferred for grazing *- Cattle Unit equivalent to a 450kg steer or a dry cow.


Land system	Description	Pastoral potential (ha/CU*)	% station
Merbla	Alluvial plains with clayey partly gilgaied soils, below gabbro hills supporting chenopod shrublands and tussock grasslands.	<35	16.1
Steer	Gravelly alluvial plains supporting chenopod shrublands.	42 - 63	3.8
Cunyu	Calcrete platforms, intervening drainage floors and channels and minor alluvial plains, supporting acacia shrublands, occasional casuarina woodlands and minor halophytic shrublands.	70 - 98	2.3
Naluthanna	Rough hills, tor fields and slopes of gabbro above lower stony plains with gilgaied drainage floors supporting mixed acacia shrublands with sparse halophytes.	70 - 98	6.4
Nubev	Gently undulating stony plains, minor limonitic low rises and drainage floors supporting mulga and halophytic shrublands.	70 - 98	1.5
Monk	Hardpan plains with occasional sandy banks supporting mulga tall shrublands and wanderie grasses.	105 - 133	10.4
Nallex	Gently undulating stony plains supporting acacia tall shrublands and chenopod low shrublands.	140 - 203	8.9

Diet selection - fDNA metabarcoding

The fDNA revealed that as many as 85 organisational taxonomic units ('Zotu), with possible 707 different plant species were eaten by cattle (Appendix 3). The most common plant species found in the faeces are listed in Figure 5. The highest sequence read numbers were recorded with Zotu1 (clade containing *Eremophila*), Zotu5 and 13 (clades containing *Sida*), Zotu6 (i.e. clade containing

Maireana), Zotu16 (clade containing *Indigofera*), and Zotu18 (clade containing *Acacia*). There was variation in individual cattle plant selection, but most cattle appeared to have consumed *Maireana* and *Eremophila species*.

Figure 5. Presence and distribution of plant species fDNA (number of sequence reads) and live weight gain (LWG) ranking in 30 cattle collected at Station A station in Apr/May 2023. Only sequence reads that were > 1000 shown. LWG ranking (g/d) - High>400; Medium 250 - 400; Low 100 - 249; Very low<100



Animal productivity

In Apr/May 2023 mustering, the LW was recorded in 49 animals. The LW was ranging from 121 to 376 (Figure 6a). Amongst 43 animals where LWG could be calculated (i.e. seen both in 2022 and 2023), there were 13 that lost weight since last weighing and 30 that gained weight (Figure 6b).

Figure 6a. LW of 49 cattle that still had Ceres tags by mustering Apr/May 2023






Figure 6b. The LWG in 43 cattle in period May 2022 – May 2023



Correlating animal location/movement with diet and animal productivity

The link between location and growth performance was assessed in 25 Ceres-tagged cattle. It was revealed that it varied in different land systems, ranging from 0.095 kg/d in Nubev to 0.389 kg/d in Naluthanna (Figure 7).

Figure 7. The LWG of individual cattle (n=25) at the preferred land systems from May 2022 to April 2023.



It appears that the growth performance was also affected by diet selection, as documented in Figure 5. Animals that had preference for species in Zotu 959, Zotu3, Zotu5 , and Zotu 41 seem to had better LWG to those that consumed Zotu1 or Zotu8 plant species.

Georeferencing of animals and alignment to vegetation characteristics.

One of the objectives of the research was to identify if georeferencing of animals could yield further information on the patterns of consumption of key vegetation characteristics. The federation of the georeferencing data series (collected using ceres ear tags) and the f DNA /vegetation data series was undertaken. Clear identification of animals with differing performance was observed with individual animals residing within different vegetation zones (Figures 8 and 9).




Figure 8 Location of cattle x fDNA federated data. Cattle #2 and #3 as high live weight gain and high abundance of Zotu3 (Abutilon clade) and Zotu4 (mixed species clade); Cattle #24 that had medium live weight gain and high abundance of Zotu1 (Eremophila clade) and Zotu8 (mixed species clade); and Cattle #26 and #32 that had very low live weight gain and high abundance of Zotu 1 only (Eremophila clade).




Figure 9 Kernel boundaries of cattle x fDNA federated data. Cattle #2 and #3 as high live weight gain and high abundance of Zotu3 (Abutilon clade) and Zotu4 (mixed species clade); Cattle #24 that had medium live weight gain and high abundance of Zotu1 (Eremophila clade) and Zotu8 (mixed species clade); and Cattle #26 and #32 that had very low live weight gain and high abundance of Zotu 1 only (Eremophila clade).

Furthermore, there were identifiable differences in animal performance and residence location. Animals 2 and 3 within zotu 3 were gaining weight rapidly, and almost exclusively within the Abutilon vegetation characteristic zone compared to Animals 24, 26 and 32 who were georeferenced to areas with vegetation dominated by Eremophila (a low digestibility but potentially high antimethanogenic vegetation clade) or mixed Eremophila vegetation clades. Further information on these issues is presented in P.PSH.1262. These data are a preliminary analysis of an extensive and extremely complex data set. However, the analysis has demonstrated that the data collected and approaches to data federation are sufficiently granular to determine the environments where individual cattle are grazing, what those cattle may consume and how long those cattle reside within those specific vegetation environments.

Discussion

The current study demonstrated that it may be possible to use a combination of geo-referencing and molecular investigation tools to start understanding the diet selection in the WA rangeland cattle. It was obvious that cattle had some preferred grazing areas. In these areas, they appeared to have grazed a wide range of plants, but the most frequent and abundant in the faeces were likely to be *Maireana*, *Acacia*, *Indigofera*, *Eremophila* and *Sida*. The productivity (LW and LWG) was quite variable amongst individual cattle, with up to three-fold differences in these values between different land systems. It appears that the land system Naluthanna that was supporting mixed acacia shrublands and halophytes resulted in the highest LWG. The value of this finding is knowing of what animals are eating

(the most) in the landscape, as well as what may be supporting their productivity. Based on this study, we may only speculate that plants identified in the fDNA and in the landscape may be supporting animal production, as more studies are needed to confirm this observation.

The study also identified some obstacles and limitations to such approach. Firstly, there were some issues related with Ceres tags - while 60 cattle were initially fitted with tags, 11 tags either fell off, missing or were not functioning. Then, from these 49 cattle, only 43 had both LW data from 2022 and 2023, so that LWG could be assessed, and only 30 cattle were sampled for fDNA at mustering in 2023. Due to excessive flooding in the area, the cattle could not be accessed for the good part of 2022, so the faecal sampling and weighing occurred only once instead of four times. As the timing and limitation of access to the cattle throughout the year resulted in a single 'spot' sample taken for fDNA analyses, the diet selection captured in such sample may only be reflective of the diet selection near to mustering time, so further studies are needed with more frequent sampling throughout the year. Whilst the fDNA metabarcoding was conducted against a more-relevant WA rangeland library compared to a global library, it was initially developed for Pilbara species. Although there are many species that overlap between northern and central WA rangelands, this Pilbara database may still be missing key species present in the central WA rangelands.

The preliminary analysis of the federated georeferenced data, fDNA and vegetation characteristics provided the first insights into the relationship between animal performance and foraging theory within West Australian rangeland environments. Foraging theory in free ranging ruminants identifies that animal performance is closely linked to energy maximisation theory. Energy maximisation is defined as the animal of known genotype consuming a mix of species that optimise rate of digestion and hence yield of energy from the feed base. The key issue is that the animal must maintain intakes of feeds that result in synchronisation of nutrients for microbial processing as well as ensuring uptake of products of digestion. The limited data analysis demonstrated clear foraging environments for individuals with two animals geolocating in vegetation environments that seem to drive high levels of animal performance (northern location, abutilon zotu fDNA reference sites) whereas others maintain a georeferenced within rangeland vegetation zones that may be of lower nutrient density (reflected by lower animal performance). The reasons for these differences in georeferencing are poorly understood as all animals have an opportunity to access vegetation zones (free ranging) of differing nutrient profiles.

Furthermore, the data may provide opportunities to design strategic supplementation strategies. Strategic supplementation in rangeland environments should not be designed for the average animal within an average vegetation environment. If animals georeference to distinct locations within the rangeland environment, an understanding of animal performance and targets for the farming system are critical to designing efficient use of supplements. Supplementation is costly, and therefore, animals that maintain high levels of performance within a rangeland environment without supplements should not be the target to delivery of this technology. Arguably strategic supplementation should be used in environments where cattle are observed frequently (high rates of residence) but do not perform well (i.e. achieve the minimum benchmark for animal performance but no more, or do not achieve the benchmark performance). In the specific example presented in this report, the target for strategic supplementation to gain maximum benefit would be within zotu 3 and 4 that is where animal performance is suboptimal compared to animals within zotu 1 and 8 (above benchmark performance). By understanding the nutritive value of feeds within these different vegetation environments, supplements can be designed and tailored to the nutritional requirements benchmarked against animal groups within certain vegetation environments. This targeted approach would result in more animals achieving benchmark weights for transitioning to backgrounding and/or lot feeding. Inevitably, a producer cannot tailor supplementation strategy to all individuals but can achieve better use of supplements across broad groups of animals within known vegetation zones.

This project also has an important link to the work being conducted on antimethanogenic plants (P.PSH.1262) and the use of virtual fencing technologies (P.PSH.1306). By understanding the location and movement patterns of animals across the rangeland environments, this project identified vegetation clades that may have nutritional attributes that drive higher levels of animal performance. That information can be used by the pastoralist in the development of management plans for rangeland environments including the use of virtual fencing technologies and supplementation strategies. Virtual fencing technologies provide an opportunity to manage animals across extent rangeland areas. P.PSH.1306 has demonstrated, using large mobs of cattle, that animals can be maintained in specific areas of rangelands thereby maximising the use of the feed base. The work in our current project (P.PSH.1245) provides granular information on grazing preference and areas that are being actively grazed. Integration of virtual fencing would reduce the risk of landscape and biodiversity degradation through overstocking and over grazing of vegetation resources.

Key findings: Implementing Ceres tags and Mapipedia can be a useful tool for developing geo-referenced 'heat maps' of rangeland cattle and preferred land systems for grazing. There was a wide range in animal performances that can potentially be linked broadly to land systems and, through fDNA metabarcoding, to plant species diversity and the nutritional value that the diversity brings to the diet.

5. Conclusion

5.1 Key findings

This project developed and validated a novel approach for feedbase mapping in the WA Rangelands. Firstly, the geo-referencing via Ceres tags and GPS collars is identified as a valuable tool to reveal where animals are grazing. In conjunction with other data such as land systems and profiles, revealed a preference for certain locations and vegetation profiles and allowed identification of preferred grazing areas. It was also identified that there is a wide choice of plants in the rangelands with favourable NV, which in turn can advance productivity of beef cattle. The fDNA metabarcoding had proven to be good tool for more accurate identification of what cattle actually graze. Some preliminary data on diet selection are obtained and potential grazing palate in the central and northern WA Rangelands revealed. The fDNA analysis also enabled discovery of many additional species that have not been reported earlier using classical approaches. Using a local laboratory and database proven to have the advantage over using a foreign laboratory and a global database, in the sense that it allowed more precise identification of plants that are present in Australia and the region. It appears that preliminary data on animal LWG can be linked to their grazing location and plants to identify what is driving the productivity.

In this project, we also identified some obstacles and limitations to such approach. For example, initial objective was to obtain real-time data on individual weights using Walk-Over-Weighing (WoW). This approach was reliant on cattle accessing water points. However, the unusual rain pattern generated excess water on the targeted property, which discouraged cattle to access water points, and hence not going through WoW scales. In future studies, other approaches may be considered such as offering supplementary feed or attractant to ensure that cattle are using WoW.

There was also limitation on how many cattle were able to be accessed to collect faecal samples from. This was reliant on weather conditions and consequently only limited number of samples were collected and analysed. The weather conditions and ability to access rangeland cattle therefore needs to be factored in when planning further trials. Another limitation to fDNA metabarcoding is that it is just a spot sample – i.e. it does not represent whole range of plants that animal may have grazed, as the plant DNA may be digested well before sample was taken. From DNA analyses perspective, there were also some inaccuracies when the foreign laboratory was used, as they mis-detected plant species not present in Australia. The local plant database, whilst having relevant plant species was also limited, as it only contains around 200 sequences of plant species, mostly from Pilbara.

5.2 Benefits to industry

Quantifying diet composition and selection of beef cattle in WA Rangelands can help in decision-making by pastoralists to achieve desired objectives through predicting outcomes of grazing strategies and identifying key plant species to guide management. Such information is critical because individual cattle consume different quantities of nutrients and animal nutritional needs fluctuate. Diet composition of animals in extensive and spatially heterogeneous environments is difficult to identify because it is hard to locate animals, plant composition varies spatial and temporal and finding, identifying and analysing plants is difficult, complex, and time consuming.

The information that is collected will be used to form guidelines for grazing management decisions and improved production. A better understanding of these factors will guide management decisions and potentially improve production. The potential production benefits from improved management decisions include improved performance of individual animals, a reduced 'tail' in the spread of liveweights, higher reproduction rates, and greater predictability in achieving target live weights.

6. Future research and recommendations

This project, (and other outputs from the WA BEEFLINKS) will form the basis for developing grazing management practices that improve production, deliver a more consistent supply of animals, and improve rangeland health. One of the main outcomes from this project is the development of guidelines for improving grazing management to better utilise the mosaic of feed sources available to increase productivity and sustainability of beef production. These guidelines will be a blueprint for all northern WA beef producers to enhance the supply chain at its origin and, in combination with the BeefLinks – Growing WA backgrounding through adoption project, improve the efficiency down the entire supply chain from north-south. The fDNA metabarcoding is a unique method still in development, but as technology advances and its use becomes more widespread, the procedure will become standardized and accurate. Ideally, a pilot study for each new application should be conducted to ensure that the sampling and analysis design is appropriate to detect the target.

Database quality and imprecise taxonomic resolution are issues yet to be overcome. Future work should focus on i) improving the reference library with more sequences of plant species relevant for the whole WA rangelands; ii) understanding limitations of the method, i.e. that multiple species may be attributed to the same Zotu; iii) greater verification of fDNA metabarcoding results using field-derived plant composition data, especially in rangeland settings where plant diversity can be intrinsically high.

While the current project was guided to collect and analyse plants based on the pre-existing information and knowledge, the later geo-referencing detection of preferred grazing area and looking at fDNA revealed what cattle may have grazed. Once these are defined, it is possible to go back to these areas and look specifically for plant species identified with the later approach. To extend further on this, those specific plants can be collected and analysed for NV. This can then truly match land system x plant species x nutritive values x what animals are grazing x productivity.

These preliminary findings also need to be validated further over a range of land conditions and plant biodiversity profiles by collecting more plants that can be added to the local fDNA database and matching with fDNA grazing profiles and changes in LW. It may be useful to access faecal samples from other trials in free ranging or setup more controlled experiments where the diet profile is well known and defined. It is also possible to setup a highly controlled digestion experiments (i.e. *in vitro* in batch, Rusitec, dual fermentors, or *in vivo* in fistulated animals) to examine how enteric digestion may affect recovery of different plant species DNA.

Apart from the animal productivity, the current approach of geo-referencing coupled with fDNA metabarcoding may be a very useful tool in for estimating rangeland land and pasture condition, and help in managing land use. For example, it can monitor the grazing pressure on vulnerable plant communities. In rehabilitation efforts, by exposing land to some cattle and then follow-up with their fDNA metabarcoding analysis, it can potentially allow early detection of plants that are not visible in the landscape or low in biomass.

7. References

- Boyer, F., Mercier, C., Bonin, A., Le Bras, Y., Taberlet, P., & Coissac, E. (2016). obitools: a unix-inspired software package for DNA metabarcoding. *Mol Ecol Resour*, 16(1), 176-182. <https://doi.org/10.1111/1755-0998.12428> [doi]
- Calderón-Sanou, I., Münkemüller, T., Boyer, F., Zinger, L., & Thuiller, W. (2020). From environmental DNA sequences to ecological conclusions: How strong is the influence of methodological choices? *Journal of Biogeography*, 47(1), 193-206. <https://doi.org/https://doi.org/10.1111/jbi.13681>
- CEAC. (2023). *Station A Station ESRM Plan*.
- Craine, J. M., Towne, E. G., Miller, M., & Fierer, N. (2015). Climatic warming and the future of bison as grazers. *Scientific Reports*, 5(1), 16738. <https://doi.org/10.1038/srep16738>
- Dixon, R., & Coates, D. (2009). Review: Near Infrared Spectroscopy of Faeces to Evaluate the Nutrition and Physiology of Herbivores. *Journal of Near Infrared Spectroscopy*, 17(1), 1-31. <https://opg.optica.org/jnirs/abstract.cfm?URI=jnirs-17-1-1>
- DPIRD. (2019). *The Western Australian Beef Industry - Pastoral Production Systems*. <https://www.agric.wa.gov.au/export-services/western-australian-beef-industry?page=0%2C2>
- Durmic, Z., Hutton, P., Revell, D. K., Emms, J., Hughes, S., & Vercoe, P. E. (2010). *In vitro* fermentative traits of Australian woody perennial plant species that may be considered as potential sources of feed for grazing ruminants. *Animal Feed Science and Technology*, 160, 98–109. <https://doi.org/10.1016/j.anifeedsci.2010.07.006>
- Durmic, Z., Moate, P. J., Jacobs, J. L., Vadhanabhuti, J., & Vercoe, P. E. (2016). *In vitro* fermentability and methane production of some alternative forages in Australia. *Animal Production Science*, 56(3), 641-645. <https://doi.org/10.1071/AN15486>
- Durmic, Z., Ramirez-Restrepo, C. A., Gardiner, C., O'Neill, C. J., Hussein, E., & Vercoe, P. E. (2017). Differences in the nutrient concentrations, *in vitro* methanogenic potential and other fermentative traits of tropical grasses and legumes for beef production systems in northern

- Australia. *Journal of the Science of Food and Agriculture*, 97, 4075-4086.
<https://doi.org/10.1002/jsfa.8274>
- Durmic, Z., Stevens, J. C., Merritt, D. J., & Erickson, T. E. (2021). *Evaluating native Pilbara plants for alternative seed-use supplies whilst providing ecosystem support and meeting mine rehabilitation targets* NRM in the Rangelands Conference - Shaping our Future, Longreach, Queensland.
- FFI. (2011). *Perennial forage shrubs providing profitable and sustainable grazing : key practical findings from the Enrich project*.
- Ficetola, G. F., Coissac, E., Zundel, S., Riaz, T., Shehzad, W., Bessière, J., Taberlet, P., & Pompanon, F. (2010). An In silico approach for the evaluation of DNA barcodes. *BMC Genomics*, 11(1), 434.
<https://doi.org/10.1186/1471-2164-11-434>
- Gwatirisa, C., Mudereri, B. T., Chitata, T., Mukanga, C., Ngwenya, M. M., Muzvondiwa, J. V., Mugandani, R., & Sungirai, M. (2022). Microhabitat and patch selection detection from GPS tracking collars of semi-free ranging Mashona cattle within a semi-arid environment. *Livestock Science*, 261, 104963. <https://doi.org/10.1016/j.livsci.2022.104963>
- Kartzinel, T. R., Chen, P. A., Coverdale, T. C., Erickson, D. L., Kress, W. J., Kuzmina, M. L., Rubenstein, D. I., Wang, W., & Pringle, R. M. (2015). DNA metabarcoding illuminates dietary niche partitioning by African large herbivores. *Proc Natl Acad Sci U S A*, 112(26), 8019-8024.
<https://doi.org/1503283112> [pii]
- 201503283 [pii]
- 10.1073/pnas.1503283112 [doi]
- Pompanon, F., Deagle, B. E., Symondson, W. O. C., Brown, D. S., Jarman, S. N., & Taberlet, P. (2012). Who is eating what: diet assessment using next generation sequencing. *Molecular Ecology*, 21(8), 1931-1950. <https://doi.org/10.1111/j.1365-294X.2011.05403.x>
- Scasta, J. D., Jorns, T., Derner, J. D., Lake, S., Augustine, D. J., Windh, J. L., & Smith, T. L. (2019). Validation of DNA metabarcoding of fecal samples using cattle fed known rations. *Animal Feed Science and Technology*, 255, 114219.
<https://doi.org/10.1016/j.anifeedsci.2019.114219>
- Taberlet, P., Bonin, A., Zinger, L., & Coissac, E. (2018). *Environmental DNA: For Biodiversity Research and Monitoring*. Oxford University Press.
<https://doi.org/10.1093/oso/9780198767220.001.0001>
- Taberlet, P., Coissac, E., Hajibabaei, M., & Rieseberg, L. H. (2012). Environmental DNA. *Molecular Ecology*, 21(8), 1789-1793. <https://doi.org/10.1111/j.1365-294X.2012.05542.x>
- Valentini, A., Pompanon, F., & Taberlet, P. (2009). DNA barcoding for ecologists. *Trends Ecol Evol*, 24(2), 110-117. [https://doi.org/S0169-5347\(08\)00344-3](https://doi.org/S0169-5347(08)00344-3) [pii]
- 10.1016/j.tree.2008.09.011 [doi]
- Van Soest, P. J., Robertson, J. B., & Lewis, B. A. (1991). Methods for dietary fiber, neutral detergent fiber, and nonstarch polysaccharides in relation to animal nutrition. *Journal Animal Science*, 74, 3583-3597.

Appendix

Appendix 1 – CN30-DietID DATABASE

Attached as Excel

Appendix 2 – Methodologies used for fDNA metabarcoding

**Appendix 3 – Presence and distribution of plant species fDNA (number of sequence reads)
in cattle sampled at Station A station in April/May 2023**

Attached as Excel

