

How to identify a phosphorus deficiency

It's not uncommon for beef producers to be unaware that a phosphorus problem exists in their herd. If stock are gaining weight and body condition is maintained once the wet season has started, there can be a misconception that stock are getting the nutrients they need from pasture – which isn't always the case.

Here are some ways to identify a phosphorus deficiency:

CLINICAL SIGNS

While clinical signs such as 'peg leg' – where cattle develop an arched body, staggering gait and brittle bones – are a sure indicator of an acute P deficiency, serious production issues could be occurring long before the deficiency is identified in the herd.

SOIL

Soil maps can provide a general indication of phosphorus deficiency. Soil P is difficult to use for diagnosis at the paddock level where there can be marked differences in soil types within a paddock.

Careful soil sampling which captures the variability across large paddocks is required to obtain a full appreciation of soil P status (Table 1). This may be expensive if multiple samples are needed. Soil Colwell P levels of >8 mg/kg suggest that the pastures contain sufficient P.

Table 1. Phosphorus (P) status of beef cattle and indicative soil P levels

| P status | Acute | Deficient | Marginal | Adequate |
|-----------------------------|-------|-----------|----------|----------|
| Colwell P levels PB (mg/kg) | <4 | 4–6 | 6–8 | >8 |

COLLECTING SAMPLES

Blood



Blood

Measuring the 'plasma inorganic phosphorus' (PiP) levels of cattle that have not been fed P supplements for at least two weeks over the wet season is the best method to determine if the pastures are providing enough P (the P-Screen Test).

To collect a sample, randomly select 20–25 head (growing animals, heifers in early-mid pregnancy or lactating first-calf cows) grazing the test paddock. Sample late in the growing season when both dietary protein and energy are considered adequate.

In continuously mated herds, introduction of growing steers (prior to the wet season) or pregnant heifers can improve the confidence in the diagnosis. Because the PiP test is the most accurate indicator of dietary P, it can also be used to assess the effectiveness of supplementation programs. However, testing supplemented animals will not establish whether or not you need to supplement in the first place.

Table 2 below shows the best estimates of P status in growing and breeder cattle for a range of plasma inorganic phosphorus levels and diet dry matter digestibility.

Table 2. P status in cattle for a range of plasma inorganic phosphorus (PiP) levels and diet dry matter (DM) digestibility

| Category | Diet DM digestibility | | |
|--------------------------------------------------------------------------------------|-----------------------|----------------------------|-----------------------------------------------------------------------|
| | >60% (Good pastures) | 55–60% (Moderate pastures) | <54% (Low quality pasture providing maintenance requirements or less) |
| Growing cattle and breeders not-lactating and up to the last two months of pregnancy | | | |
| <1.0 | Acutely deficient | Acutely deficient | Deficient |
| 1.0–1.5 | Acutely deficient | Deficient | Marginal |
| 1.5–2.0 | Deficient | Marginal | Adequate |
| Breeders in the last two months of pregnancy and early to mid-lactation | | | |
| <1.0 | Acutely deficient | Acutely deficient | Deficient |
| 1.0–1.3 | Acutely deficient | Deficient | Deficient |
| 1.3–1.5 | Deficient | Marginal | Marginal |

If PiP is >2.0 mmol/L, cattle are generally expected to have adequate P intake. If dry matter digestibility is less than 50%, no growth response is expected – although there may be bone P replenishment.



What's the impact of a phosphorus deficiency in beef herds?

Phosphorus supplementation in acutely deficient pastures will increase feed intake over the growing season when both energy and protein are adequate and P has become the primary limiting nutrient. Because P supplementation of deficient cattle significantly increases feed intake, cattle numbers must be adjusted to maintain sustainable long-term carrying capacity.

Phosphorus deficiency results in poor appetite and feed intake, poor growth, high breeder mortality rates, reduced fertility and milk production, bone breakage and, in severe cases, bone deformities and peg leg.

Maintaining phosphorus reserves is particularly important for breeder cows when they're lactating and/or are in the last trimester of pregnancy. Extra phosphorus is required for milk production and for the bones and tissues of the developing foetus. When there's insufficient phosphorus in the diet, it's mobilised from the bones, which can increase bone fragility and injuries in stock being handled.

Faecal

The main pathway of P excretion from the body is faeces. Chemical analysis of faecal P can be a guide if P intake is very low or high, but not in marginal deficiency situations. However, P supplementation must be removed for at least two weeks prior to sampling for an accurate measure, which could be highly impractical.

Pastures

Pasture samples can provide an indication of phosphorus levels, but the animals' daily grazing selection could mean the samples may not accurately represent the actual plants consumed.

Aim to evaluate the P status of a paddock in normal seasons – avoid exceptionally good or poor years.

Clinical signs of P deficiency such as peg leg are most evident in dry/drought years but the productivity gains apply to all seasons.

Bones

Bones are extremely useful indicators of an existing phosphorus deficiency; however, the tests require either a bone biopsy or an X-Ray – neither of which are cost-effective or practical for commercial properties.

Supplementing P-deficient beef herds

All classes of cattle, whether they be steers, heifers or breeders, will benefit from phosphorus supplementation; however, the response of the cattle depends on the severity of deficiency, class of animal and diet quality. The greatest benefits from supplementation on deficient pastures are achieved over the wet season when grass is green and stock are gaining weight.

Breeders in late pregnancy or lactating breeders grazing dry season pastures in phosphorus-deficient country still have high P requirements.

For steers and heifers, providing additional supplementary phosphorus to animals on a protein-deficient diet during the dry season can be detrimental if nitrogen (protein) is not included in the supplement. During the dry season, protein becomes the primary limiting nutrient and feeding a 'P only' supplement will not improve performance but can be detrimental and even induce bone fragility.

Supplementation guide

The quality of inorganic phosphorus supplements can vary depending on the source and how they've been made. It's important to ask for an analysis of the supplement you choose and focus on the bioavailability of the phosphorus (what's available to the animal) rather than the total amount.

As a general guide, mono-calcium phosphate (MCP) has the highest bioavailability, followed closely by mono-dicalcium phosphate (MDCP) and dicalcium phosphate (DCP). MDCP is a blend of MCP and DCP but isn't always in the same proportions, which is why quality and pricing may vary.

If you use DCP, ensure it's hydrous DCP so that it's soluble in water – otherwise, it may be totally ineffective. Tricalcium phosphate (TCP) and rock phosphate aren't recommended as supplements for cattle due to the extremely low bioavailability of phosphorus.

Requirements

Unlike urea supplementation, there's no toxicity issue associated with phosphorus supplementation (unless urea is part of the supplement). The issue is not so much 'over-supplementing', but meeting the challenge of getting adequate phosphorus into cattle diets over the wet season when there's more opportunity for increased weight gains.

Table 3: Recommended supplementary P intakes for several classes of cattle and live weight changes

| Class of animal, live weight (kg) and weight gain (kg) | Supplementary P intake required (g/head/day) | | | |
|--------------------------------------------------------|----------------------------------------------|---------------------|-------------|------------|
| | Target weight gain (kg/day) | Acutely P deficient | P deficient | Marginal P |
| Wet season | | | | |
| Pregnant breeders (400kg) | 0 | 5 | 3 | 1 |
| | 0.6 | 9 | 6 | 2 |
| Steers and heifers (200kg) | 0 | 2 | 1 | 0 |
| | 0.6 | 6 | 4 | 1 |
| Steers and heifers (400kg) | 0 | 3 | 2 | 1 |
| | 0.6 | 7 | 5 | 2 |
| Dry season | | | | |
| Late-pregnant or lactating cows (400kg) | -0.3 | 7 | 5 | 2 |
| | 0 | 9 | 6 | 2 |

^A Live weight change as limited by pasture protein and energy intake when diet P is adequate

Top tips

- Be aware that some sources of inorganic phosphorus may contain traces of heavy metals that may accumulate over an extended feeding period and could potentially show up in meat in rare and extreme cases.
- Consider the best supplement system for your situation e.g. loose mixes (covered lick sheds or exposed bulka bags), blocks or water medication.
- Excessive amounts of calcium should be avoided.
- Water medication systems can be impractical where alternative surface water is available.
- In phosphorus-deficient high rainfall regions with sown pastures, phosphorus fertilisers are often more cost-effective than supplementing stock directly. In these situations, fertilisers increase pasture yields and diet quality.

Are there any other considerations?

There are some other important factors to consider when compiling a phosphorus supplementation plan for northern beef herds.



Cropping: Phosphorus deficiency can occur in places where it's not expected to be a problem. Cereal cropping, especially when not accompanied by appropriate fertiliser application, has caused soil nutrient rundown in some regions. In these situations, cattle performance may be improved with phosphorus supplements. A serum phosphorus test is the best way to diagnose the problem.



Assessment of individual paddocks: Where marked soil variation occurs within properties, it's recommended to assess every major paddock when developing a comprehensive P supplementation program for the enterprise. If the assessment is performed on young grower or breeder cattle that have not received P supplement for at least two weeks prior to sampling and are on a positive plane of nutrition, this will most likely be a 'one-off' test unless phosphorus fertiliser is being applied.



Botulism: Phosphorus supplementation does not reduce the need to vaccinate against botulism. It will reduce the risk by avoiding the bone-chewing behavior that can lead to botulism.



Cost of supplementation: The cost benefits will vary depending on the level of deficiency being encountered. A 2018 trial in the NT suggested a cost of around \$20/head/year to provide adequate phosphorus for breeders, although cost varies depending on the current price of phosphorus. Other costs such as distribution, freight and storage costs will change between properties and this also needs to be factored into the economic assessment. In this trial, the benefits of supplementation outweighed the cost:

- 30kg increase in annual heifer growth rates
- >30kg increase in weaner weights
- >100kg increase in breeder cow weights
- at least a 7% reduction in mortality rates
- at least a 30% increase in re-conception rates.

Conduct your own cost-benefit analysis depending on the severity of the deficiency in your pastures.



More information

Resources

Phosphorus management of beef cattle in northern Australia (2012). Meat & Livestock Australia

Quigley, S., Poppi, D., and Schatz T. (2015). *Validation and demonstration of a diagnostic tool for phosphorus status of beef cattle*. MLA Final Report, Meat & Livestock Australia

Schatz, T. and McCosker, K. (2018). *Phosphorus supplementation of Brahman heifers in phosphorus deficient country in the NT*. Proceedings Australian Society of Animal Production, Category Animal Nutrition, Ref 2695

Dixon, R., Anderson, S., Kidd, L., and Fletcher, M. (2020). *Management of phosphorus nutrition of beef cattle grazing seasonally dry rangelands: a review*. Animal Production Science, CSIRO

Dixon, R., Fletcher, M., Anderson, A., Kidd, L., Benvenuti, M., Mayer, B., McNeill, D. and Goodwin, L. (2020). *Improved management of cattle phosphorus status through applied physiology*. MLA Final Report, Meat & Livestock Australia

Contact


Nigel Tomkins
E: ntomkins@mla.com.au

- Several trials have demonstrated the benefits of phosphorus intake:**
- ▲ 15–30% increase in **weaning rates**
 - ▲ Up to 130kg increases in **breeder cow live weights**
 - ▲ 6–12% increase in **birth weights**
 - ▼ 15–50% decrease in **breeder mortality rates**

What has the research revealed?

Recent MLA breeder cow supplementation trials have provided a better understanding of phosphorus's role in northern beef herds, including when and where to provide the supplement cost-effectively.

Figure 2: The impact of phosphorus supplementation on feed intake and live weight gains on pen-fed steers



Care is taken to ensure the accuracy of the information contained in this publication. However, MLA cannot accept responsibility for the accuracy or completeness of the information or opinions contained in the publication. You should make your own enquiries before making decisions concerning your interests. MLA accepts no liability for any losses incurred if you rely solely on this publication and excludes all liability as a result of reliance by any person on such information or advice.

Apart from any use permitted under the Copyright Act 1968, all rights are expressly reserved. Requests for further authorisation should be directed to the Content Manager, PO Box 1961, North Sydney, NSW 2059 or info@mla.com.au. © Meat & Livestock Australia 2020 ABN 39 081 678 364. Published in April 2021.

MLA acknowledges the matching funds provided by the Australian Government to support the research and development detailed in this publication.

PO Box 1961
North Sydney NSW 2059
P: 02 9463 9333
E: info@mla.com.au
mla.com.au



Why do cattle need phosphorus?

A guide for northern beef producers



- P¹⁵ Phosphorus (P) is:**
- a major contributor to muscle and bone health
 - essential for metabolism
 - particularly important for improving cattle growth, health and milk production.

The need for phosphorus

For many regions in northern Australia, phosphorus (P) deficiency is a serious nutritional issue for beef herds and can cause major losses in productivity and profitability.

Cattle need phosphorus for almost every vital function of the body. It's used for:

- building bones and teeth
- metabolising fat, carbohydrates and protein
- producing milk
- increasing feed intake.

Deficiencies often arise in tropical production systems because most northern soils have lower available soil P compared with southern Australia. This means there's often insufficient phosphorus in the pasture to meet animal requirements.

The cattle that have the highest phosphorus requirements are growing stock, late-pregnant heifers and cows, and lactating cows.

Figure 1: How phosphorus is used in beef cattle

